

MATH 1020 WORKSHEET 10.3
Polar Coordinates and Polar Graphs

Polar coordinates require the basic transformation equations

$$\begin{aligned}x &= r \cos \theta \\r^2 &= x^2 + y^2\end{aligned}$$

$$\begin{aligned}y &= r \sin \theta \\ \tan \theta &= y/x\end{aligned}$$

Given the polar point $(-1, 5\pi/4)$, find the corresponding Cartesian coordinates for the point.

Solution. We apply the transformation equations to determine the x and y coordinates of the point.

$$\begin{aligned}x &= -1 \cdot \cos\left(\frac{5\pi}{4}\right) & y &= -1 \cdot \sin\left(\frac{5\pi}{4}\right) \\x &= -1 \left(\frac{-1}{\sqrt{2}}\right) & y &= -1 \left(\frac{-1}{\sqrt{2}}\right) \\x &= \frac{1}{\sqrt{2}} & y &= \frac{1}{\sqrt{2}}\end{aligned}$$

Thus, writing the answer as a point we have

$$\boxed{\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)}$$

Convert the polar equation $r = 4 \sin \theta$ to a Cartesian form and sketch its graph.

Solution. Transforming to a polar equation, it is best to start by replacing any trigonometric functions in the equation before regrouping and then replacing any r terms.

$$\begin{aligned}r &= 4 \sin \theta \\&= 4 \left(\frac{y}{r}\right) \\r^2 &= 4y \\x^2 + y^2 &= 4y\end{aligned}$$

From here, the easiest way to sketch the equation is to bring the $4y$ over to the LHS and complete the square so that the Cartesian equation is in a familiar form.

$$\begin{aligned}x^2 + y^2 - 4y &= 0 \\x^2 + (y^2 - 4y + \underline{\quad}) &= 0 + \underline{\quad} \\x^2 + (y^2 - 4y + \underline{4}) &= 0 + \underline{4} \\x^2 + (y - 2)^2 &= 4\end{aligned}$$

One recognizes that we have the equation of a circle. Thus

$$\boxed{x^2 + y^2 = 4y \text{ is a circle with center at } (0, 2) \text{ and radius } r = 2.}$$

Convert the Cartesian equation $xy = 4$ to a polar form.

Solution. Converting a Cartesian equation into polar form is done by substituting the transformation equations for $x = r \cos \theta$ and $y = r \sin \theta$ into the given equation.

$$\boxed{r^2 \cos \theta \sin \theta = 4}.$$

Find the slope of the tangent line to the polar curve $r = 2 - \sin \theta$ at $\theta = \frac{\pi}{3}$.

Solution. To determine slope of the tangent line to the polar curve, we must first convert the polar equations into parametric form using the transformation equations. For parametric form we have

$$\begin{aligned}x &= r \cos \theta & y &= r \sin \theta \\x &= (2 - \sin \theta) \cos \theta & y &= (2 - \sin \theta) \sin \theta \\x &= 2 \cos \theta - \cos \theta \sin \theta & y &= 2 \sin \theta - \sin^2 \theta\end{aligned}$$

Now we need to determine $\frac{dx}{d\theta}$ and $\frac{dy}{d\theta}$.

$$\frac{dx}{d\theta} = -2 \sin \theta - \cos^2 \theta + \sin^2 \theta \quad \frac{dy}{d\theta} = 2 \cos \theta - 2 \sin \theta \cos \theta$$

Substituting into the parametric formula for slope we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{dy/d\theta}{dx/d\theta} \\&= \frac{2 \cos \theta - 2 \sin \theta \cos \theta}{-2 \sin \theta - \cos^2 \theta + \sin^2 \theta}\end{aligned}$$

Now we can evaluate at $\theta = \pi/3$

$$\begin{aligned}\left. \frac{dy}{dx} \right|_{\theta=\pi/3} &= \left. \frac{2 \cos \theta - 2 \sin \theta \cos \theta}{-2 \sin \theta - \cos^2 \theta + \sin^2 \theta} \right|_{\theta=\pi/3} \\&= \frac{2 \cos(\pi/3) - 2 \sin(\pi/3) \cos(\pi/3)}{-2 \sin(\pi/3) - \cos^2(\pi/3) + \sin^2(\pi/3)} \\&= \frac{2 \left(\frac{1}{2}\right) - 2 \left(\frac{\sqrt{3}}{2}\right) \left(\frac{1}{2}\right)}{-2 \left(\frac{\sqrt{3}}{2}\right) - \left(\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} \\&= \frac{1 - \frac{\sqrt{3}}{2}}{-\sqrt{3} - \frac{1}{4} + \frac{3}{4}} \\&= \frac{1 - \frac{\sqrt{3}}{2}}{-\sqrt{3} + \frac{1}{2}}\end{aligned}$$

Thus the slope of the tangent line to the polar curve $r = 2 - \sin \theta$ at $\theta = \frac{\pi}{3}$ is

$$\boxed{\left. \frac{dy}{dx} \right|_{\theta=\pi/3} = \frac{2 - \sqrt{3}}{-2\sqrt{3} + 1}}.$$