

Exercise 1: Find the domain for the given rational function.

$$f(x) = \frac{3x^2 + x}{x^2 + 4}$$

Exercise 2: Find the domain for the given rational function.

$$f(x) = \frac{3x^2}{2x^2 - 5x - 3}$$

Answers

Exercise 1: Find the domain for the given rational function.

$$f(x) = \frac{3x^2 + x}{x^2 + 4}$$

To determine if we have any domain restrictions, set the denominator equal to zero and solve for x.

$$x^2 + 4 = 0$$

$$x^2 = -4$$

x^2 will never be equal to a negative number so there are no domain restrictions. The domain for this rational function is all real numbers.

$$D = (-\infty, \infty) \text{ -- interval notation}$$

$$\text{or } D = \{x \mid x \in \mathbb{R}\} \text{ -- set notation}$$

Exercise 2: Find the domain for the given rational function.

$$f(x) = \frac{3x^2}{2x^2 - 5x - 3}$$

Set denominator equal to zero and solve for x

$$2x^2 - 5x - 3 = 0$$

$$(2x + 1)(x - 3) = 0$$

Set each factor equal to zero

$$2x + 1 = 0 \text{ or } x - 3 = 0$$

$$x = -\frac{1}{2} \text{ or } x = 3$$

Our domain restrictions are that x cannot be equal to $-1/2$ or 3, so our domain will be all real numbers except for these two.

$$D = \left(-\infty, -\frac{1}{2}\right) \cup \left(-\frac{1}{2}, 3\right) \cup (3, \infty)$$

$$\text{or } D = \left\{x \mid x \neq -\frac{1}{2} \text{ or } 3\right\}$$

Exercise 3: Find all vertical and horizontal asymptotes (if any) for the given rational function.

$$f(x) = \frac{3x+4}{x-5}$$

Answers

Exercise 3: Find all vertical and horizontal asymptotes (if any) for the given rational function.

$$f(x) = \frac{3x+4}{x-5}$$

To find the vertical asymptote, set the denominator equal to zero and solve for x.

$$x - 5 = 0$$

$$x = 5$$

To find the horizontal asymptote, compare the degree power of the numerator and denominator.

numerator is $3x^1 + 4$ with a degree of 1

denominator is $x^1 - 5$ with a degree of 1

Since the degree powers are the same the horizontal asymptote will be equal to the ratio of the leading coefficients in the numerator and denominator.

$$f(x) = \frac{3x+4}{1x-5}$$

$$y = \frac{3}{1}$$

$$y = 3$$

The asymptotes are:

vertical: $x = 5$

horizontal: $y = 3$

Exercise 4: Find all vertical and horizontal asymptotes (if any) for the given rational function.

$$f(x) = \frac{x^2 + 4x + 4}{x^3 - x^2 - 6x}$$

Exercise 5: Determine the x and y intercepts, vertical and horizontal asymptotes, symmetry, and any additional points needed to graph the given rational function.

$$f(x) = \frac{2x^2 - 18}{-3x^2 - 12x - 12}$$

Answers

Exercise 5: Determine the x and y intercepts, vertical and horizontal asymptotes, symmetry, and any additional points needed to graph the given rational function.

$$f(x) = \frac{2x^2 - 18}{-3x^2 - 12x - 12}$$

Set $f(x) = 0$ and solve for x to find the x-intercepts

$$0 = \frac{2x^2 - 18}{-3x^2 - 12x - 12}$$

$$0 = 2x^2 - 18$$

$$18 = 2x^2$$

$$9 = x^2$$

$$\pm 3 = x$$

x-intercepts are (-3,0) and (3, 0)

Substitute 0 for x to find the y-intercept

$$f(x) = \frac{2x^2 - 18}{-3x^2 - 12x - 12}$$

$$f(0) = \frac{2(0)^2 - 18}{-3(0)^2 - 12(0) - 12}$$

$$= \frac{-18}{-12}$$

$$= \frac{3}{2}$$

y-intercept is (0, 3/2)

Answers

Substitute $-f(x)$ for $f(x)$ to test for symmetry to the x-axis

$$\begin{aligned}f(x) &= \frac{2x^2 - 18}{-3x^2 - 12x - 12} \\-f(x) &= -\frac{2x^2 - 18}{-3x^2 - 12x - 12} \\&= \frac{-2x^2 + 18}{-3x^2 - 12x - 12}\end{aligned}$$

$-f(x)$ is not the same as $f(x)$ so the function is not symmetric to the x-axis.

Substitute $-x$ for x and $-f(x)$ for $f(x)$ to test for symmetry to the origin

$$\begin{aligned}f(x) &= \frac{2x^2 - 18}{-3x^2 - 12x - 12} \\-f(-x) &= -\frac{2(-x)^2 - 18}{-3(-x)^2 - 12(-x) - 12} \\&= -\frac{2x^2 - 18}{-3x^2 + 12x - 12} \\&= \frac{-2x^2 + 18}{-3x^2 + 12x - 12}\end{aligned}$$

$-f(-x)$ is not the same as $f(x)$ so the function is not symmetric to the origin.

Evaluate $f(x)$ at -7 and 7 to get a couple more points for the graph.

$$\begin{aligned}f(x) &= \frac{2x^2 - 18}{-3x^2 - 12x - 12} & f(x) &= \frac{2x^2 - 18}{-3x^2 - 12x - 12} \\f(-7) &= \frac{2(-7)^2 - 18}{-3(-7)^2 - 12(-7) - 12} & f(7) &= \frac{2(7)^2 - 18}{-3(7)^2 - 12(7) - 12} \\&= \frac{80}{-75} & &= \frac{80}{-243} \\&\approx -1.07 & &\approx -0.33\end{aligned}$$

Answers

Exercise 4: Find all vertical and horizontal asymptotes (if any) for the given rational function.

$$f(x) = \frac{x^2 + 4x + 4}{x^3 - x^2 - 6x}$$

Vertical asymptotes

$$x^3 - x^2 - 6x = 0$$

$$x(x^2 - x - 6) = 0$$

$$x(x - 3)(x + 2) = 0$$

$$x = 0 \text{ or } x - 3 = 0 \text{ or } x + 2 = 0$$

$$x = 0 \text{ or } x = 3 \text{ or } x = -2$$

Horizontal asymptotes

numerator is $x^2 + 4x + 4$ with a degree of 2

denominator is $x^3 - x^2 - 6x$ with a degree of 3

Since the degree power of the denominator is one more than the numerator, the horizontal asymptote will be the x-axis.

$$y = 0$$

The asymptotes are:

vertical: $x = -2$, $x = 0$, and $x = 3$

horizontal: $y = 0$