




# MATRICES AND MATRIX OPERATIONS

**MULTIPLYING MATRICES** – We can multiply only matrices where the first matrix has the number of columns same as the number of rows of the second matrix. And new matrix AB will have same number of rows as the first matrix, and same number of columns as the second matrix. The next drawing will help you to understand.



**Note:**  $A \times B = AB$  and  $B \times A = BA$ , but when we are multiplying matrices AB isn't the same as BA.

**!!!  $AB \neq BA$**

Can we multiply next matrices?

$$1) \quad A = \begin{bmatrix} 1 & 2 & 4 \\ 2 & 6 & 0 \end{bmatrix} \quad B = \begin{bmatrix} 4 & 0 & 2 \\ 1 & 1 & 3 \\ 5 & 2 & 7 \\ 8 & 9 & 1 \end{bmatrix} \quad A \quad B \quad = \quad AB \\
 \begin{array}{c}
 \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\
 2 \times 3 \quad 4 \times 3 \quad = \quad ?
 \end{array}$$

Matrix A has size 2x3 (2 rows and 3 columns), and matrix B has size 4x3 (4 rows and 3 columns). The number of columns of the matrix A is 3, and the number of the rows of the matrix B is 4, as these numbers are not the same we CAN'T multiply these two matrices.

$$2) \quad A = \begin{bmatrix} 1 & 2 & 4 \\ 2 & 6 & 0 \end{bmatrix} \quad B = \begin{bmatrix} 4 & 1 & 4 & 3 \\ 0 & -1 & 3 & 1 \\ 2 & 7 & 5 & 2 \end{bmatrix} \quad A \quad B \quad = \quad AB \\
 \begin{array}{c}
 \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\
 2 \times 3 \quad 3 \times 4 \quad = \quad 2 \times 4
 \end{array}$$

Matrix A has size 2x3 (2 rows and 3 columns), and matrix B has size 3x4 (3 rows and 4 columns). The number of columns of the matrix A is 3, and the number of the rows of the matrix B is 3, as these numbers are the same we CAN multiply these two matrices.

Now, when we know that we can multiply A and B, we need to see what is going to be the size of the product matrix AB. Number of the rows of the matrix AB is equal to the number of the rows of the matrix A, it is 2. Number of the column of the matrix AB is equal to the number of the columns of the matrix B, it is 4. Thus, the size of the matrix AB is 2 x 4.



# MATRICES AND MATRIX OPERATIONS

**TRANSPOSE OF THE MATRIX** - It is a new matrix that we get when rows and columns of the matrix A change places, transpose of A is denoted by  $A^T$ . If matrix A has size  $m \times n$ , then matrix  $A^T$  will have size  $n \times m$ , because we have changed row and columns.

$$\text{a) } A = \begin{bmatrix} 2 & 1 & 5 \\ 3 & 4 & 6 \end{bmatrix} \quad A^T = ? \quad \text{b) } B = \begin{bmatrix} 7 & 2 & 4 \\ -2 & 3 & 5 \\ 1 & -8 & 9 \end{bmatrix} \quad B^T = ?$$

To find  $A^T$  rows and columns of matrix A need to change places. We will put the first row [2 1 5] as a first column of  $A^T$ , and the second row [3 4 6] as second column of the  $A^T$ . Do the same for  $B^T$ .

$$\text{a) } A^T = \begin{bmatrix} 2 & 3 \\ 1 & 4 \\ 5 & 6 \end{bmatrix} \quad \text{b) } B^T = \begin{bmatrix} 7 & -2 & 1 \\ 2 & 3 & -8 \\ 4 & 5 & 9 \end{bmatrix}$$

**Note:** If matrix is squared and  $A = A^T$  we say that it is SYMMETRIC MATRIX.

$$A = \begin{bmatrix} 5 & -2 & -1 \\ -2 & 4 & 7 \\ -1 & 7 & 6 \end{bmatrix} \quad \text{if we change places for rows and columns we will get } A^T = \begin{bmatrix} 5 & -2 & -1 \\ -2 & 4 & 7 \\ -1 & 7 & 6 \end{bmatrix}$$

From here we can see  $A = A^T$ , thus this matrix is symmetric.

**Rules for transpose** (if the sizes of matrices are such that stated operations can be performed):

- $(A^T)^T = A$
- $(kA)^T = kA^T$ , k is scalar (number)
- $(A \pm B)^T = A^T \pm B^T$
- $(AB)^T = B^T A^T$

**MINORS OF MATRIX** - In this handout we will only cover minors for  $3 \times 3$  matrices, but similarly it can be calculated for any squared matrix

For doing this we need to know determinate of the matrix  $2 \times 2$ .

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \det(A) = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

If A is a squared matrix, then the minor, denoted by  $M_{ij}$ , of element  $a_{ij}$  is the determinate of submatrix that remains after the  $i$ th row and  $j$ th column are deleted from A.

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3} \quad \det(A) = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

# MATRICES AND MATRIX OPERATIONS

If we want  $M_{11}$ , since that is the minor that correspond to element  $a_{11}$ , we are going to cover row 1 and column 1, everything that is left we will write in the same order in our minor.

$$\begin{vmatrix} \cancel{a_{11}} & \cancel{a_{12}} & \cancel{a_{13}} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \quad M_{11} = \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix}$$

Now we are going to do the same thing for every minor.

$$M_{12} = \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} \quad M_{13} = \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

$$M_{21} = \begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix} \quad M_{22} = \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} \quad M_{23} = \begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix}$$

$$M_{31} = \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix} \quad M_{32} = \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix} \quad M_{33} = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$$

**COFACTORS** - the number  $C_{ij} = (-1)^{i+j} \times M_{ij}$  is the cofactor of element  $a_{ij}$

**ADJOINT OF THE MATRIX (ADJUGATE)**

$$\mathbf{M} = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix} = \begin{bmatrix} +M_{11} & -M_{12} & +M_{13} \\ -M_{21} & +M_{22} & -M_{23} \\ +M_{31} & -M_{32} & +M_{33} \end{bmatrix} = \begin{bmatrix} + \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} & - \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} & + \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} \\ - \begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix} & + \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} & - \begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix} \\ + \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix} & - \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix} & + \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} \end{bmatrix}$$

$$\mathbf{adj}(A) = \mathbf{M}^T$$

Note that signs (+ or -) are easy to remember where to put the right one. Start with + and then -, +, -, +, etc.

**INVERSE OF MATRIX** -If A is the square matrix and B is the same size of A. If matrix B can be find such that  $AB = BA = I$ , then A is said to be invertible (nonsingular, if  $\det(A) \neq 0$ ), and B is called an inverse of A. If there is no such matrix B, then A is not invertible (singular, if  $\det(A) = 0$ ).

Notation for the inverse of matrix A is  $A^{-1}$ . If B is inverse of A, then  $B = A^{-1}$ .

$$AB = BA = I \text{ or } AA^{-1} = A^{-1}A = I$$

Inverse of matrix A (for all squared matrices) can be found using this formula:  $A^{-1} = \frac{1}{\det(A)} \mathbf{adj}(A)$ .

Inverse of 2x2 matrix  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  is  $A^{-1} = \frac{1}{\det(A)} \mathbf{adj}(A) = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$ , if and only if  $\det(A) \neq 0$ .

## MATRICES AND MATRIX OPERATIONS

$$a) A = \begin{bmatrix} 6 & 1 \\ 5 & 2 \end{bmatrix}, A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{6 \times 2 - 1 \times 5} \begin{bmatrix} 2 & -1 \\ -5 & 6 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 2 & -1 \\ -5 & 6 \end{bmatrix} = \begin{bmatrix} \frac{2}{7} & -\frac{1}{7} \\ -\frac{5}{7} & \frac{6}{7} \end{bmatrix}$$

$$b) A = \begin{bmatrix} 1 & 5 & 0 \\ 0 & 3 & 2 \\ 1 & 0 & 2 \end{bmatrix}_{3 \times 3}, A^{-1} = ?$$

$$M_{11} = \begin{vmatrix} 3 & 2 \\ 0 & 2 \end{vmatrix} = 3 \times 2 - 0 \times 2 = 6 \qquad M_{12} = \begin{vmatrix} 0 & 2 \\ 1 & 2 \end{vmatrix} = -2 \qquad M_{13} = \begin{vmatrix} 0 & 3 \\ 1 & 0 \end{vmatrix} = -3$$

$$M_{21} = \begin{vmatrix} 5 & 0 \\ 0 & 2 \end{vmatrix} = 10 \qquad M_{22} = \begin{vmatrix} 1 & 0 \\ 1 & 2 \end{vmatrix} = 2 \qquad M_{23} = \begin{vmatrix} 1 & 5 \\ 1 & 0 \end{vmatrix} = -5$$

$$M_{31} = \begin{vmatrix} 5 & 0 \\ 3 & 2 \end{vmatrix} = 10 \qquad M_{32} = \begin{vmatrix} 1 & 0 \\ 0 & 2 \end{vmatrix} = 2 \qquad M_{33} = \begin{vmatrix} 1 & 5 \\ 0 & 3 \end{vmatrix} = 3$$

$$M = \begin{bmatrix} +6 & -(-2) & +(-3) \\ -10 & +2 & -(-5) \\ +10 & -2 & +3 \end{bmatrix} = \begin{bmatrix} 6 & 2 & -3 \\ -10 & 2 & 5 \\ 10 & -2 & 3 \end{bmatrix} \quad adj(A) = M^T = \begin{bmatrix} 6 & -10 & 10 \\ 2 & 2 & -2 \\ -3 & 5 & 3 \end{bmatrix}$$

Help - Determinant of only 3x3 matrix can be find using Sarrus' rule. We write first two columns of the determinate to the right of the determinant (in that order). Then we are adding the products of the diagonals, going from the top to bottom (dashed lines), and subtract products of the diagonals going from the bottom to the top (solid lines).

$$det(A) = \begin{vmatrix} 1 & 5 & 0 & 1 & 5 \\ 0 & 3 & 2 & 0 & 3 \\ 1 & 0 & 2 & 1 & 0 \end{vmatrix} = 1 \times 3 \times 2 + 5 \times 2 \times 1 + 0 \times 0 \times 0 - 1 \times 3 \times 0 - 0 \times 2 \times 1 - 2 \times 0 \times 5 = 6 + 10 + 0 - 0 - 0 - 0 = 16$$

$$\text{Finally, } A^{-1} = \frac{1}{det(A)} adj(A) = \frac{1}{16} \begin{bmatrix} 6 & -10 & 10 \\ 2 & 2 & -2 \\ -3 & 5 & 3 \end{bmatrix} = \begin{bmatrix} \frac{6}{16} & \frac{-10}{16} & \frac{10}{16} \\ \frac{2}{16} & \frac{2}{16} & \frac{-2}{16} \\ \frac{-3}{16} & \frac{5}{16} & \frac{3}{16} \end{bmatrix} = \begin{bmatrix} \frac{3}{8} & \frac{-5}{8} & \frac{5}{8} \\ \frac{1}{8} & \frac{1}{8} & \frac{-1}{8} \\ \frac{-3}{16} & \frac{5}{16} & \frac{3}{16} \end{bmatrix}$$

**Rules for inverse** (if A is invertible):

- $(A^{-1})^T = (A^T)^{-1}$
- $(A^{-1})^{-1} = A$
- $(AB)^{-1} = B^{-1}A^{-1}$
- $(kA)^{-1} = k^{-1}A^{-1} = \frac{1}{k}A^{-1}$ , k is nonzero scalar
- $(A_1A_2 \dots A_n)^{-1} = A_n^{-1} \dots A_2^{-1}A_1^{-1}$

**References:** The following work were referred to during the creation of this handout: *Elementary Linear Algebra, Application Version, 11<sup>th</sup> Ed., Howard Anthon, Chriss Roress; and* <http://www.matematiranje.in.rs>