

Alternate Definition of Derivative ... Facts 1

The Limit Definition of Derivative

The **derivative** of a function f with respect to the variable x is the function f' whose value at x is

f' prime of x ←

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h},$$

provided the limit exists.

→ you can take the derivative

Anywhere that the derivative exists, we say that the function is *differentiable*.

Alternate Definition of Derivative ... Facts 1

Thus, the derivative f' is a function that gives the slope of the function f at any point.

Other Notation used with derivatives (we will use most of these so remember them).

if equation is $y =$
the derivative will be $\frac{dy}{dx}$ or y'

Alternate Definition of Derivative ... Facts 1

Example 1: Use the definition of derivative to find $f'(x)$. $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

a) $f(x) = 3x + 2$

$$f(x+h) = 3(x+h) + 2 = 3x + 3h + 2$$

$$\lim_{h \rightarrow 0} \frac{3x + 3h + 2 - 3x - 2}{h} \rightarrow \lim_{h \rightarrow 0} \frac{3h}{h} = 3 \text{ so, } f'(x) = 3$$

Alternate Definition of Derivative ... Facts 1

b) $f(x) = x^3 + x^2$

$$f(x+h) = (x+h)^3 + (x+h)^2 = x^3 + 3x^2h + 3xh^2 + h^3 + x^2 + 2xh + h^2$$

$$f(x) = x^3 + x^2$$

$$\lim_{h \rightarrow 0} \frac{3x^2h + 3xh^2 + h^3 + 2xh + h^2}{h} \Rightarrow \boxed{3x^2 + 2x = f'(x)}$$

Alternate Definition of Derivative ... Facts 1

Modified Form of the Limit Definition of the Derivative

The **numeric value** of the derivative of a function f at the point $(c, f(c))$ is given as

$$f'(c) = \lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h},$$

provided the limit exists.

Alternate Definition of Derivative ... Facts 1

If you are asked to find the derivative at a point, you have two options: you can find the formula first then plug in the point, or you can plug in the point in the beginning. However, if you need to find the derivative at multiple points, I suggest finding the formula first to save yourself some time.

Example 2: If $f(x) = 5x^2 + 2x$, find $f'(2)$.

$$\begin{aligned} f(x+h) &= 5(x+h)^2 + 2(x+h) \\ &= 5(x^2 + 2xh + h^2) + 2x + 2h \\ &= 5x^2 + 10xh + 5h^2 + 2x + 2h \\ \lim_{h \rightarrow 0} \frac{5x^2 + 10xh + 5h^2 + 2x + 2h - 5x^2 - 2x}{h} \end{aligned}$$

$$\lim_{h \rightarrow 0} \frac{10xh + 5h^2 + 2h}{h}$$

$$\lim_{h \rightarrow 0} 10x + 5h + 2$$

$$f'(x) = 10x + 2$$

$$f'(2) = 10(2) + 2 = 22$$


Alternate Definition of Derivative ... Facts 1

Alternative Definition of Derivative

An alternative definition of the derivative of f at point c is

$$f'(c) = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c},$$

provided the limit exists.



What this alternative definition allows us to do is to examine the behavior of a function as x approaches c from the left or the right. The limit exists (and thus the derivative) as long as the left and right limits exist and are equal.

Alternate Definition of Derivative ... Facts 1

Example 3: Use the alternative definition of the derivative to find the derivative of $f(x) = \frac{1}{\sqrt{x}}$ at $x = 4$

$$\lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4}$$

$$f(4) = \frac{1}{\sqrt{4}} = \frac{1}{2}$$

$$\lim_{x \rightarrow 4} \frac{\frac{1}{\sqrt{x}} - \frac{1}{2}}{x - 4}$$

$$\lim_{x \rightarrow 4} \frac{\frac{2 - \sqrt{x}}{2\sqrt{x}} \cdot \frac{2 + \sqrt{x}}{2 + \sqrt{x}}}{x - 4}$$

$$\lim_{x \rightarrow 4} \frac{4 - x - 1}{2\sqrt{x}(2 + \sqrt{x})(x - 4)}$$

$$\frac{-1}{2\sqrt{4}(2 + \sqrt{4})} = \frac{-1}{2(2) \cdot 4} = \boxed{\frac{-1}{16} = f'(4)}$$

Alternate Definition of Derivative ... Facts 1

Example 4: If $f(x) = x^2 - 4x$, answer the following.

a) Find the average rate of change of $f(x)$ over the interval $[-1, 2]$.

x-values

$$f(-1) = (-1)^2 - 4(-1) = 1 + 4 = 5$$
$$f(2) = (2)^2 - 4(2) = 4 - 8 = -4$$
$$\frac{f(2) - f(-1)}{2 - (-1)} = \frac{-4 - 5}{2 + 1} = \frac{-9}{3} = \boxed{-3}$$

b) Find the instantaneous rate of change of $f(x)$ at the point where $x = 2$.

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \quad \text{or} \quad f'(2) = \lim_{x \rightarrow 2} \frac{f(x) - f(2)}{x - 2}$$

$$f'(x) = 2x - 4 \quad \text{so...} \quad f'(2) = 2(2) - 4 = \boxed{0} \quad \text{😊}$$



Alternate Definition of Derivative ... Facts 1

Relationships between the graphs of f and f'

Since a derivative at any point is equivalent to the slope of the function at that point, we can graph the derivative given a function and estimate what the original function looks like when we are given the graph of the derivative.

Example 5: Given the graph of f , sketch the graph of the derivative on the same set of axes.

f'




Alternate Definition of Derivative ... Facts 1

Example 6: Given the graph of f' and $f(0) = 2$, sketch the graph of the function f on the same set of axes.

$f'(x) = -1$ if $x < 1$
 so $f(x) = -x + c$
↓
unknown

$f'(x) = 2$ if $x \geq 1$
 $f(x) = 2x + c$
↓
unknown



Why is it necessary to know $f(0) = 2$?

I need to "where" (so y-int is helpful)

Alternate Definition of Derivative ... Facts 1

Example 7: Suppose the graph below is the graph of the derivative of h .

a) What is the value of $h'(0)$? What does this tell us about $h(x)$?


$$h'(0) = 4$$

↓ at $x=0$, $h(x)$ is increasing at a rate of 4

b) Using the graph of $h'(x)$, how can we determine when the graph of $h(x)$ is going up? How about going down?

↙ when $h'(x)$ is positive (above the x-axis)

↘ when $h(x)$ is negative (below x-axis)



c) The graph of $h'(x)$ crosses the x-axis at $x = 2$ and $x = -2$. Describe the behavior of the graph of $h(x)$ at these points.

↪ this is where the slope is 0, so the graph will change direction