


Tangent and Normal Lines ... Facts 2

Tangents and Normals

Equation of a Tangent Line

The derivative at a point $x = a$, denoted $f'(a)$, is the instantaneous rate of change at that point. Geometrically, $f'(a)$ gives us the **slope** of the tangent line at the point $x = a$.



Recall: A **tangent line** is a line that "just touches" a curve at a specific point without intersecting it.

To find the **equation of the tangent line** we need its slope and a point on the line. Given the function $f(x)$ and the point $(a, f(a))$ we can find the equation of the tangent line using the slope equation.

$$m = \frac{f(x) - f(a)}{x - a}$$

Since $f'(a)$ gives us the slope of the tangent line at the point $x = a$, we have

$$f'(a) = \frac{f(x) - f(a)}{x - a}$$

As such, the equation of the tangent line at $x = a$ can be expressed as:


$$f(x) - f(a) = f'(a)(x - a)$$

Tangent and Normal Lines ... Facts 2

Equation of a Normal Line

The **normal line** is defined as the line that is perpendicular to the tangent line at the point of tangency. Knowing this, we can find the equation of the normal line at $x = a$ by taking the negative inverse of the slope of the tangent line equation.

Thus, if $f'(a)$ is the slope of the tangent line at $x = a$. The negative inverse is $\frac{-1}{f'(a)}$.



As such, the equation of the normal line at $x = a$ can be expressed as:

$$f(x) - f(a) = \frac{-1}{f'(a)}(x - a)$$

Tangent and Normal Lines ... Facts 2

Example 1: Find the equation of the tangent and normal lines of the function $f(x) = \sqrt{2x - 1}$ at the point (5, 3).

Solution:

a) Equation of the Tangent Line.

<p>Step 1: Find the slope of the function by solving for its first derivative.</p>	$f(x) = \sqrt{2x - 1}$ $f(x) = (2x - 1)^{\frac{1}{2}}$ $f'(x) = \frac{1}{2}(2x - 1)^{-\frac{1}{2}}(2)$ $f'(x) = \frac{1}{\sqrt{2x - 1}}$
<p>Step 2: Knowing $f'(a)$, solve for the slope of the tangent at $a = 5$.</p>	$f'(5) = \frac{1}{\sqrt{2(5) - 1}}$ $f'(5) = \frac{1}{3}$
<p>Step 3: Solve for $f(a)$.</p>	$f(5) = \sqrt{2(5) - 1}$ $f(5) = 3$
<p>Step 4: Substitute found values into the equation of a tangent line.</p>	$f(x) - f(a) = f'(a)(x - a)$ $f(x) - 3 = \frac{1}{3}(x - 5)$

Tangent and Normal Lines ... Facts 2

b) Equation of the Normal Line.

Step 1: Find the slope of the normal line $\frac{-1}{f'(a)}$.	Since $f'(5) = \frac{1}{3}$, then $\frac{-1}{f'(a)} = -3$
Step 2: Given the equation of a tangent line, swap slopes.	$f(x) - f(a) = \frac{-1}{f'(a)}(x - a)$ $f(x) - 3 = -3(x - 5)$

Tangent and Normal Lines ... Facts 2

Example 2: Find the equation of the tangent and normal lines of the function $f(x) = (x^2 - 1)^3$ at the point $(2, 27)$.

Solution:

a) Equation of the Tangent Line.

Step 1: Find the slope of the function by solving for its first derivative.	$f(x) = (x^2 - 1)^3$ $f'(x) = 3(x^2 - 1)^2(2x)$ $f'(x) = 6x(x^2 - 1)^2$
Step 2: Knowing $f'(a)$, solve for the slope of the tangent at $a = 2$.	$f'(2) = 6(2)((2)^2 - 1)^2$ $f'(2) = 108$
Step 3: Solve for $f(a)$.	$f(2) = ((2)^2 - 1)^3$ $f(2) = 27$
Step 4: Substitute found values into the equation of a tangent line.	$f(x) - f(a) = f'(a)(x - a)$ $f(x) - 27 = 108(x - 2)$

b) Equation of the Normal Line.

Step 1: Find the slope of the normal line $\frac{-1}{f'(a)}$.	Since $f'(2) = 108$, then $\frac{-1}{f'(a)} = -\frac{1}{108}$
Step 2: Given the equation of a tangent line, swap slopes.	$f(x) - f(a) = \frac{-1}{f'(a)}(x - a)$ $f(x) - 27 = -\frac{1}{108}(x - 2)$

Tangent and Normal Lines ... Facts 2

Exercises:

1. Find the equation for the normal and tangent lines for $f(x)$ at the specified points.

a) $f(x) = e^x$ at $(0,1)$

b) $f(x) = 2x^3 - 3x + 7$ at $(1,6)$

c) $f(x) = \frac{1}{x^2}$ at $(-1,1)$

d) $f(x) = x\cos(x)$ at $(0,0)$

e) $f(x) = xe^x$ at $(0, 0)$

Tangent and Normal Lines ... Facts 2

Answers

1. a) Tangent: $y - 1 = x$, Normal: $y - 1 = -x$
- b) Tangent: $y - 6 = 3(x - 1)$, Normal: $y - 6 = -\frac{1}{3}(x - 1)$
- c) Tangent: $y - 1 = 2(x + 1)$, Normal: $y - 1 = -\frac{1}{2}(x + 1)$
- d) Tangent: $y = x$, Normal: $y = -x$
- e) Tangent: $y = x$, Normal: $y = -x$