

Inverse Trig Rule ... Set 2

Calculus

Ch. 5.6 Inverse Trig Derivatives

Classwork Worksheet

THEOREM 5.16 Derivatives of Inverse Trigonometric Functions

Let u be a differentiable function of x .

$$\begin{aligned} \frac{d}{dx} [\arcsin u] &= \frac{u'}{\sqrt{1-u^2}} & \frac{d}{dx} [\arccos u] &= \frac{-u'}{\sqrt{1-u^2}} \\ \frac{d}{dx} [\arctan u] &= \frac{u'}{1+u^2} & \frac{d}{dx} [\operatorname{arccot} u] &= \frac{-u'}{1+u^2} \\ \frac{d}{dx} [\operatorname{arcsec} u] &= \frac{u'}{|u|\sqrt{u^2-1}} & \frac{d}{dx} [\operatorname{arccsc} u] &= \frac{-u'}{|u|\sqrt{u^2-1}} \end{aligned}$$

Evaluating an Expression In Exercises 21–24, evaluate each expression without using a calculator. (*Hint:* See Example 3.)

21. (a) $\sin\left(\arctan \frac{3}{4}\right)$

22. (a) $\tan\left(\arccos \frac{\sqrt{2}}{2}\right)$

23. (a) $\cot\left[\arcsin\left(-\frac{1}{2}\right)\right]$

24. (a) $\sec\left[\arctan\left(-\frac{3}{5}\right)\right]$

Simplifying an Expression Using a Right Triangle In Exercises 25–32, write the expression in algebraic form. (*Hint:* Sketch a right triangle, as demonstrated in Example 3.)

25. $\cos(\arcsin 2x)$

26. $\sec(\arctan 4x)$

29. $\tan\left(\operatorname{arcsec} \frac{x}{3}\right)$

31. $\csc\left(\arctan \frac{x}{\sqrt{2}}\right)$

Inverse Trig Rule ... Set 2

Answers

Inverse Trig Derivatives

THEOREM 5.16 Derivatives of Inverse Trigonometric Functions

Let u be a differentiable function of x .

$$\frac{d}{dx} [\arcsin u] = \frac{u'}{\sqrt{1-u^2}} \quad \frac{d}{dx} [\arccos u] = \frac{-u'}{\sqrt{1-u^2}}$$


$$\frac{d}{dx} [\arctan u] = \frac{u'}{1+u^2} \quad \frac{d}{dx} [\operatorname{arccot} u] = \frac{-u'}{1+u^2}$$

$$\frac{d}{dx} [\operatorname{arcsec} u] = \frac{u'}{|u|\sqrt{u^2-1}} \quad \frac{d}{dx} [\operatorname{arccsc} u] = \frac{-u'}{|u|\sqrt{u^2-1}}$$

Evaluating an Expression In Exercises 21–24, evaluate each expression without using a calculator. (*Hint:* See Example 3.)


21. (a) $\sin\left(\arctan \frac{3}{4}\right)$

$$= \boxed{\frac{3}{5}}$$




22. (a) $\tan\left(\arccos \frac{\sqrt{2}}{2}\right)$

$$= \frac{\sqrt{2}}{\sqrt{2}} = \boxed{1}$$



23. (a) $\cot\left[\arcsin\left(-\frac{1}{2}\right)\right]$


$$= \frac{\sqrt{3}}{-1} = \boxed{-\sqrt{3}}$$



$[-\pi/2, \pi/2]$

24. (a) $\sec\left[\arctan\left(-\frac{3}{5}\right)\right]$

$$= \frac{\sqrt{34}}{5}$$




$\tan \theta = -\frac{3}{5}$

$[-\pi/2, \pi/2]$

Simplifying an Expression Using a Right Triangle In Exercises 25–32, write the expression in algebraic form. (*Hint:* Sketch a right triangle, as demonstrated in Example 3.)


25. $\cos(\arcsin 2x)$

$$= \frac{\sqrt{1-4x^2}}{1} = \boxed{\sqrt{1-4x^2}}$$




26. $\sec(\arctan 4x)$

$$= \frac{\sqrt{1+16x^2}}{1} = \boxed{\sqrt{1+16x^2}}$$




29. $\tan\left(\operatorname{arcsec} \frac{x}{3}\right)$

$$= \frac{\sqrt{x^2-9}}{3}$$



31. $\csc\left(\arctan \frac{x}{\sqrt{2}}\right)$

$$= \frac{\sqrt{2+x^2}}{x}$$



Inverse Trig Rule ... Set 2

Let u be a differentiable function of x .

$$\frac{d}{dx} [\arcsin u] = \frac{u'}{\sqrt{1-u^2}}$$

$$\frac{d}{dx} [\arccos u] = \frac{-u'}{\sqrt{1-u^2}}$$

$$\frac{d}{dx} [\arctan u] = \frac{u'}{1+u^2}$$

$$\frac{d}{dx} [\operatorname{arccot} u] = \frac{-u'}{1+u^2}$$

$$\frac{d}{dx} [\operatorname{arcsec} u] = \frac{u'}{|u|\sqrt{u^2-1}}$$

$$\frac{d}{dx} [\operatorname{arccsc} u] = \frac{-u'}{|u|\sqrt{u^2-1}}$$

Finding a Derivative In Exercises 39–58, find the derivative of the function.

39. $f(x) = 2 \arcsin(x - 1)$

44. $f(x) = \arctan \sqrt{x}$

46. $h(x) = x^2 \arctan 5x$

47. $h(t) = \sin(\arccos t)$

50. $y = \ln(t^2 + 4) - \frac{1}{2} \arctan \frac{t}{2}$

Inverse Trig Rule ... Set 2

Answers

THEOREM 5.16 Derivatives of Inverse Trigonometric Functions

Let u be a differentiable function of x .

$$\begin{aligned} \frac{d}{dx} [\arcsin u] &= \frac{u'}{\sqrt{1-u^2}} & \frac{d}{dx} [\arccos u] &= \frac{-u'}{\sqrt{1-u^2}} \\ \frac{d}{dx} [\arctan u] &= \frac{u'}{1+u^2} & \frac{d}{dx} [\operatorname{arccot} u] &= \frac{-u'}{1+u^2} \\ \frac{d}{dx} [\operatorname{arcsec} u] &= \frac{u'}{|u|\sqrt{u^2-1}} & \frac{d}{dx} [\operatorname{arccsc} u] &= \frac{-u'}{|u|\sqrt{u^2-1}} \end{aligned}$$

Finding a Derivative In Exercises 39–58, find the derivative of the function.

39. $f(x) = 2 \arcsin(x-1)$

$$f'(x) = 2 \cdot \left(\frac{1}{\sqrt{1-(x-1)^2}} \right) = \boxed{\frac{2}{\sqrt{1-(x-1)^2}}}$$

44. $f(x) = \arctan \sqrt{x} \quad \arctan(x^{1/2})$

$$f'(x) = \frac{\frac{1}{2}x^{-1/2}}{1+(\sqrt{x})^2} = \frac{1}{2\sqrt{x}(1+x)}$$

$$= \boxed{\frac{1}{2\sqrt{x}(1+x)}}$$

46. $h(x) = x^2 \arctan 5x$

**product rule*

$$h'(x) = \underbrace{2x}_{f'} \cdot \underbrace{\arctan(5x)}_g + \underbrace{x^2}_f \cdot \underbrace{\frac{5}{1+(5x)^2}}_{g'}$$

$$h'(x) = 2x \arctan(5x) + \frac{5x^2}{1+25x^2}$$

47. $h(t) = \sin(\arccos t)$

**chain rule*

$$h'(t) = \underbrace{\cos(\arccos t)}_{\text{out: } \sin(\)} \cdot \underbrace{-\frac{1}{\sqrt{1-t^2}}}_{\text{in: } \arccos t}$$

$$h'(t) = t \cdot \left(\frac{-1}{\sqrt{1-t^2}} \right)$$

$$h'(t) = \boxed{\frac{-t}{\sqrt{1-t^2}}}$$

50. $y = \ln(t^2+4) - \frac{1}{2} \arctan \frac{t}{2}$ ← $\frac{1}{2} \arctan\left(\frac{1}{2}t\right)$

**apply*

$$\frac{d}{dx} \ln u = \frac{u'}{u}$$

$$y' = \frac{2t}{t^2+4} - \frac{1}{2} \cdot \frac{1}{1+\left(\frac{t}{2}\right)^2}$$

$$y' = \frac{2t}{t^2+4} - \frac{1}{4\left(1+\frac{t^2}{4}\right)}$$

$$y' = \frac{2t}{t^2+4} - \frac{1}{4+t^2} = \boxed{\frac{2t-1}{t^2+4}}$$