

# Curve Sketching ... Set 1

## Curve Sketching Practice

With a partner or two and without the use of a graphing calculator, attempt to sketch the graphs of the following functions. Pertinent aspects of the graph to include (include as many as you can):

- asymptotes (vertical/horizontal)
  - domain
  - local extrema/regions of increase/decrease
  - points of inflection/concavity
  - $x$ -intercepts(?)
- 

1.  $f(x) = x^4 - 6x^2$

# Curve Sketching ... Set 1

## Answers

1

The zeros ( $x$ -intercepts) of  $f$ :

$$x^4 - 6x^2 = 0 \Rightarrow x^2(x^2 - 6) = 0 \Rightarrow x = 0, \pm\sqrt{6}.$$

The zeros of  $f'(x) = 4x^3 - 12x$ :


$$4x^3 - 12x = 0 \Rightarrow 4x(x^2 - 3) = 0 \Rightarrow x = 0, \pm\sqrt{3}.$$

$f'$  changes sign at each of these numbers since  $f'(-2) < 0$ ,  $f'(-1) > 0$ ,  $f'(1) < 0$ , and  $f'(2) > 0$ . Thus,  $f$  has relative minimums  $f(-\sqrt{3}) = -9$  and  $f(\sqrt{3}) = -9$  and a relative maximum  $f(0) = 0$ .

The zeros of  $f''(x) = 12x^2 - 12$ :

$$12x^2 - 12 = 0 \Rightarrow 12(x^2 - 1) = 0 \Rightarrow x = -1, 1.$$

$f''$  changes sign as it passes each of these numbers, since  $f''(-2) > 0$ ,  $f''(0) < 0$  and  $f''(2) > 0$ , so  $f$  has points of inflection  $(-1, -5)$  (where the graph changes from being concave upward to concave downward) and  $(1, -5)$  (concave down to concave up).



# Curve Sketching ... Set 1

## Curve Sketching Practice

With a partner or two and without the use of a graphing calculator, attempt to sketch the graphs of the following functions. Pertinent aspects of the graph to include (include as many as you can):

- asymptotes (vertical/horizontal)
  - domain
  - local extrema/regions of increase/decrease
  - points of inflection/concavity
  - $x$ -intercepts(?)
- 

2.  $f(x) = (x^2 - 1)^3$

# Curve Sketching ... Set 1

## Answers

2

The zeros of  $f$ :

$$(x^2 - 1)^3 = 0 \quad \Rightarrow \quad [(x+1)(x-1)]^3 = 0 \quad \Rightarrow \quad x = -1, 1.$$

The zeros of  $f'(x) = 6x(x^2 - 1)^2$ :


$$6x(x^2 - 1)^2 = 0 \quad \Rightarrow \quad 6x(x+1)^2(x-1)^2 = 0 \quad \Rightarrow \quad x = -1, 0, 1.$$

Since  $f'(-2) < 0$ ,  $f'(-0.5) < 0$ ,  $f'(0.5) > 0$  and  $f'(2) > 0$ ,  $f$  has a relative minimum  $f(0) = -1$ . The product rule may be used to find  $f''$ :

$$f''(x) = 6(x^2 - 1)^2 + 24x^2(x^2 - 1) = 6(x^2 - 1)[(x^2 - 1) + 4x^2] = 6(x^2 - 1)(5x^2 - 1).$$

These algebraic simplifications were carried out to factor  $f''$ , so that its zeros  $x = -1, -1/\sqrt{5}, 1/\sqrt{5}, 1$  are more easily found. Since  $f''(-2) > 0$ ,  $f''(0.5) < 0$ ,  $f''(0) > 0$ ,  $f''(0.5) < 0$  and  $f''(-2) > 0$ ,  $f$  has points of inflection at  $(-1, 0)$  (where, incidentally, the tangent line is horizontal by the fact that  $f'(-1) = 0$  and the graph goes from concave upward to concave downward), at  $(-1/\sqrt{5}, -64/125)$ , at  $(1/\sqrt{5}, -64/125)$  and at  $(1, 0)$ .

Here are graphs for the functions in problems 1 and 2 sketched using the information we gained above:



# Curve Sketching ... Set 1

## Curve Sketching Practice

With a partner or two and without the use of a graphing calculator, attempt to sketch the graphs of the following functions. Pertinent aspects of the graph to include (include as many as you can):

- asymptotes (vertical/horizontal)
  - domain
  - local extrema/regions of increase/decrease
  - points of inflection/concavity
  - $x$ -intercepts(?)
- 

3.  $f(x) = x\sqrt{x^2 + 1}$

# Curve Sketching ... Set 1

## Answers

3

The zeros of  $f$  occur only when the numerator is zero – namely, at  $x = 0$ . Finding the derivative of  $f$  is a matter for the product rule. We have

$$f'(x) = (x^2 + 1)^{1/2} + x^2(x^2 + 1)^{-1/2} = \frac{(x^2 + 1) + x^2}{\sqrt{x^2 + 1}} = \frac{2x^2 + 1}{\sqrt{x^2 + 1}}.$$


Again,  $f'$  is zero only when its numerator is zero, and since the equation  $2x^2 + 1 = 0$  has no real solutions,  $f$  is not going to have any relative extrema. Turning to the question of concavity we may apply the quotient rule to  $f'$  in the form in which it appears above, but I choose here instead to write  $f'(x) = (2x^2 + 1)(x^2 + 1)^{-1/2}$  and apply the product rule to get  $f''$ :

$$\begin{aligned} f''(x) &= 4x(x^2 + 1)^{-1/2} - x(2x^2 + 1)(x^2 + 1)^{-3/2} \\ &= \frac{4x}{\sqrt{x^2 + 1}} - \frac{2x^3 + x}{(x^2 + 1)\sqrt{x^2 + 1}} \\ &= \frac{4x(x^2 + 1) - 2x^3 - x}{(x^2 + 1)\sqrt{x^2 + 1}} \\ &= \frac{2x^3 + 3x}{(x^2 + 1)\sqrt{x^2 + 1}}. \end{aligned}$$

As with any fractional expression,  $f''$  may be zero only when its numerator is zero, so we solve

$$2x^2 + 3x = 0 \quad \Rightarrow \quad x(2x + 3) = 0 \quad \Rightarrow \quad x = 0, -\frac{3}{2}.$$

Since  $f''(-2) < 0$ ,  $f''(-1) < 0$  and  $f''(1) > 0$  we know there is one inflection point at  $(0, 0)$ , with the concavity of  $f$  changing there from downward to upward.



# Curve Sketching ... Set 1

## Curve Sketching Practice

With a partner or two and without the use of a graphing calculator, attempt to sketch the graphs of the following functions. Pertinent aspects of the graph to include (include as many as you can):

- asymptotes (vertical/horizontal)
  - domain
  - local extrema/regions of increase/decrease
  - points of inflection/concavity
  - $x$ -intercepts(?)
- 

4.  $f(x) = \frac{x}{(x-1)^2}$

# Curve Sketching ... Set 1

## Answers

4

$f$  has just one zero at  $x = 0$ . Writing  $f$  in the equivalent form  $f(x) = x(x-1)^{-2}$ , we get the derivative using the product rule (of course, the quotient rule would also have been an option):

$$f'(x) = (x-1)^{-2} - 2x(x-1)^{-3} = \frac{1}{(x-1)^2} - \frac{2x}{(x-1)^3} = \frac{(x-1) - 2x}{(x-1)^3} = \frac{-x-1}{(x-1)^3}.$$

This shows that  $f'$  has only one zero at  $x = -1$ . Like usual we should check the sign of  $f'$  on both sides of this number. What is different about this particular example is that if we check the sign of, say,  $f'(0)$ , we may **not** presume that this is the sign of  $f'$  for all numbers  $x > -1$ , rather this is true just for those numbers  $-1 < x < 1$ .  $x = 1$  is a number not actually in the domain of  $f$  — actually the site of a vertical asymptote for this function — and we have to check the sign of  $f'$  for numbers  $x > 1$  separately. Since  $f'(-2) < 0$ ,  $f'(0) > 0$  and  $f'(2) < 0$  we have the  $f$  decreases on  $-\infty < x < -1$ , reaching a local minimum  $f(-1) = -\frac{1}{4}$ , and increases on  $-1 < x < 1$  (with the values of  $f$  approaching  $+\infty$ ), and decreasing (coming down from  $+\infty$ ) for  $1 < x < \infty$ . We now glean what we can from the second derivative. Writing  $f'(x)$  in the equivalent form  $(-x-1)(x-1)^{-3}$ , we get

$$\begin{aligned} f''(x) &= -(x-1)^{-3} - 3(-x-1)(x-1)^{-4} \\ &= \frac{-1}{(x-1)^3} + \frac{3x+3}{(x-1)^4} \\ &= \frac{-(x-1) + 3x+3}{(x-1)^4} \\ &= \frac{2x+4}{(x-1)^4}. \end{aligned}$$

We determine possible points of inflection first by determining where the numerator of  $f''$  is zero. The only solution to  $2x+4=0$  is  $x=-2$ . We note that  $f''(-3) < 0$ ,  $f''(0) > 0$ ,  $f''(2) > 0$  (we check numbers from the two intervals  $-2 < x < 1$  and  $1 < x < +\infty$  separately because of the break in the domain at  $x=1$ ). Thus  $f$  is concave down on the interval  $-\infty < x < -2$ , has a point of inflection at  $(-2, -2/9)$ , and is concave up on each of the intervals  $-2 < x < 1$  and  $1 < x < +\infty$ .

